

Baryon moments in a QCD-based model*

Loyal Durand and Phuoc Ha
 Department of Physics, University of Wisconsin
 Madison, Wisconsin 53706, USA

We have analyzed the theory of the baryon magnetic moments in the (approximate) QCD setting suggested by Brambilla *et al.* By modifying their derivation of the qqq interaction and wave equation for baryons, we derived expressions for the baryon moments in terms of the underlying quark moments, including the first corrections associated with the binding of the quarks in baryons. The results, which hold in the “quenched approximation” in which the contributions of virtual quark pairs are neglected, fail to describe the measured moments, with typical errors on the order of 7%. We conclude from our analysis that the quenched approximation is at fault, and that the baryon moments give a sensitive test of that standard approximation in the lattice and analytic approaches to QCD.

1 Introduction

The magnetic moments of the stable baryons are very well measured, and their theoretical calculation gives a sensitive test of our understanding of baryon structure in quantum chromodynamics (QCD). Although the basic pattern and approximate magnitudes of the moments can be explained using the nonrelativistic quark model, the deviation of the moments from the quark-model pattern has not been explained dynamically despite many attempts. We discuss that problem here from the point of view of nonperturbative QCD. In particular, we have analyzed the theory of the moments in the approximate QCD setting developed in the work of Brambilla *et al.* [1], who used a Wilson-loop approach to study the qqq bound state problem. By modifying their derivation of the qqq interaction and wave equation, we have derived expressions for the baryon moments in terms of the underlying quark moments, including the first corrections associated with the binding of the quarks in baryons. Our results hold in the “quenched” approximation in which there are no internal quark loops embedded in the world sheet swept out by the Wilson lines joining the valence quarks, and no pairs associated with the valence lines. The corrections to the moments are

* Presented by L. Durand at the Como Conference on *Quark Confinement and the Hadron Spectrum II*, June, 1996

of relative order $\langle V \rangle / m_q$, where V is a typical component of the binding potential and m_q is a constituent quark mass, and are potentially large enough to explain the deviations of the measured moments from the quark-model values.

To test the theory, we have constructed variational wave functions for the baryons using the interactions derived by Brambilla *et al.* [1], including all spin and orbital configurations possible for $J^P = \frac{1}{2}^+$ and internal and total orbital angular momenta $L \leq 2$, and used them to calculate the moments. The effects of excited orbital contributions to the moments are negligible, as expected. The new contributions tend to cancel, and the remnants do not have the correct pattern to explain the discrepancies between theory and experiment.

The approximations underlying the Wilson-line analysis are known. Since of these, only the quenched approximation has a direct effect on the spin dependent terms with which we are dealing, we conclude that the use of the quenched approximation is responsible for the deviations of theory from experiment, and that the moment problem provides a sensitive test of this standard approximation in lattice and analytic QCD.

In a world-sheet picture, the inclusion of internal quark loops which describe meson emission and absorption by the baryon would allow the introduction of internal orbital angular momentum and spin, and would affect the moments. We are now investigating meson-loop contributions using approximations suggested by the world-sheet picture and chiral perturbation theory. In the following sections, we sketch our derivations and calculations, and justify our conclusions. A more detailed discussion will be given elsewhere.

2 Baryon moments in QCD

2.1 The problem

The simple, nonrelativistic quark model gives a qualitatively good description of the baryon moments. Under the assumption that each baryon is composed of three valence or constituent quarks in a state with all internal orbital angular momenta equal to zero, the moments are given by expectation values of the spin moment operators

$$\mu_{\text{QM}} = \sum_{\text{q}} \mu_{\text{q}} \sigma_{\text{q}}, \quad (1)$$

leading to the standard expressions

$$\mu_{\text{p}} = \frac{1}{3}(4\mu_{\text{u}} - \mu_{\text{d}}), \dots, \quad \mu_{\text{q}} = \frac{e_{\text{q}}}{2m_{\text{q}}}. \quad (2)$$

The masses in the quark moments are clearly effective masses, and can be treated as free parameters in attempting to fit the data. A least-squares fit to the measured octet moments, taken with equal weights, is shown in Table 1. The moments are given in nuclear magnetons (nm). The pattern of the signs of

the quark model moments agrees with observation, while the root-mean-square deviation of theory from experiment is 0.14 nuclear magnetons, or about 9% of the average magnitudes of the moments. Agreement at this level can be regarded as an outstanding success of the quark model, but the deviations also give a very sensitive test of baryon structure: there is presently no completely successful first-principles theory of the moments.

Table 1: Quark model fit to the magnetic moments in the baryon octet. All moments are given in nuclear magnetons.

Baryon	Experiment	Quark Model	$\Delta\mu$
p	2.793	2.728	0.065
n	-1.913	-1.819	-0.094
Σ^+	2.458 ± 0.010	2.639	-0.181
Σ^-	-1.160 ± 0.025	-0.999	-0.161
$ \Sigma^0 \rightarrow \Lambda\gamma $	1.61 ± 0.08	1.575	-0.03
Λ	-0.613 ± 0.004	-0.642	0.029
Ξ^0	-1.250 ± 0.014	-1.462	0.212
Ξ^-	-0.651 ± 0.025	-0.553	-0.098

2.2 The Wilson-loop approach to the baryon moments

Our approach to the baryon moment problem is based on the work of Brambilla *et al.* [1], who derived the interaction potential and wave equation for the valence quarks in a baryon from QCD using a Wilson-line construction. The basic idea is to construct a Green's function for the propagation of a gauge-invariant combination of quarks joined by path ordered Wilson-line factors

$$U = P \exp i g \int A_g \cdot dx, \quad (3)$$

where A_g is the color gauge field. The Wilson lines sweep out a three-sheeted world sheet of the form shown in Fig. 1 as the quarks move from their initial to their final configurations.

By making an expansion in powers of $1/m_q$ and considering only forward propagation of the quarks in time using the Foldy-Wouthuysen approximation, Brambilla *et al.* are able to derive a Hamiltonian and Schrödinger equation for the quarks, with an interaction which involves an average over the gauge field. That average is performed using the minimal surface approximation in which fluctuations in the world sheet are ignored, and the geometry is chosen to minimize the total area of the world sheet *per* Wilson, subject to the motion of the quarks. The short-distance QCD interactions are taken into account explicitly.




Figure 1: World sheet picture for the structure of a baryon.

The result of this construction is an effective Hamiltonian to be used in a semirelativistic Schrödinger equation $H\Psi = e\Psi$,

$$\begin{aligned}
H = & \sum_i \sqrt{p_i^2 + m_i^2} + \sigma(r_1 + r_2 + r_3) - \sum_{i < j} \frac{2}{3} \frac{\alpha_s}{r_{ij}} \\
& - \frac{1}{2m_1^2} \frac{\sigma}{r_1} \mathbf{s}_1 \cdot (\mathbf{r}_1 \times \mathbf{p}_1) + \frac{1}{3m_1^2} \mathbf{s}_1 \cdot [(\mathbf{r}_{12} \times \mathbf{p}_1) \frac{\alpha_s}{r_{12}^3} + (\mathbf{r}_{13} \times \mathbf{p}_1) \frac{\alpha_s}{r_{13}^3}] \\
& - \frac{2}{3} \frac{1}{m_1 m_2} \frac{\alpha_s}{r_{12}^3} \mathbf{s}_1 \cdot \mathbf{r}_{12} \times \mathbf{p}_2 - \frac{2}{3} \frac{1}{m_1 m_3} \frac{\alpha_s}{r_{13}^3} \mathbf{s}_1 \cdot \mathbf{r}_{13} \times \mathbf{p}_3 + \dots \quad (4)
\end{aligned}$$

Here $\mathbf{r}_{ij} = \mathbf{x}_i - \mathbf{x}_j$ is the separation of quarks i and j , r_i is the distance of quark i from point at which the sum $r_1 + r_2 + r_3$ is minimized, and $\mathbf{p}_i$ and $\mathbf{s}_i$ are the momentum and spin operators for quark i . The parameter σ is a “string tension” which specifies the strength of the long range confining interaction, and α_s is the strong coupling. The terms hidden in the ellipsis include tensor and spin-spin interactions which will not play a role in the analysis of corrections to the moment operator, and the terms that result from permutations of the particle labels. The full Hamiltonian is given in [1]. This Hamiltonian, including the terms omitted here, gives a good description of the baryon spectrum as shown by Carlson, Kogut, and Pandharipande [2] and by Capstick and Isgur [3], who proposed it on the basis of reasonable physical arguments, but did not give formal derivations from QCD.

The presence of the quark momenta $\mathbf{p}_i$ in the Thomas-type spin-dependent interactions in Eq. (4) suggests that new contributions to the magnetic moment operator could arise in a complete theory through the minimal substitution

$$\mathbf{p}_i \rightarrow \mathbf{p}_i - e_i \mathbf{A}_{\text{em}}(x_i), \quad (5)$$

with $\mathbf{A}_{\text{em}}(x_i)$ the electromagnetic vector potential associated with an external magnetic field. However, this point is obscured in the work of Brambilla *et al.* by the transformations they make to express the equations for the Green's function in Wilson-loop form. We have therefore repeated their derivation of the valence-quark Hamiltonian in Eq. (4), replacing the $\text{SU}(3)_c$ color gauge field gA_g by the full $\text{SU}(3)_c \times \text{U}(1)_{\text{em}}$ gauge field $gA_g + e_q A_{\text{em}}$ at the beginning. By then reorganizing the calculation of the three-quark Hamiltonian to keep $e_q A_{\text{em}}$ explicit throughout, and then expanding to first order in A_{em} with

$$\mathbf{A}_{\text{em}} = \frac{1}{2} \mathbf{B} \times \mathbf{x}_q, \quad \mathbf{B} = \text{constant}, \quad (6)$$

we can identify the modified magnetic moment operator through the relation

$$\Delta H = -\boldsymbol{\mu} \cdot \mathbf{B}. \quad (7)$$

The new moment operator, $\boldsymbol{\mu} = \boldsymbol{\mu}_{\text{QM}} + \Delta\boldsymbol{\mu}$, involves the leading corrections to the quark-model operator associated with the binding interactions. $\Delta\boldsymbol{\mu}$ can, in fact, be read off from the terms in Eq. (4) which depend on both the quark spins and momenta by making the minimal substitution in Eq. (5). For example, the term which depends on $\mathbf{s}_1 \cdot \mathbf{r}_{12} \times \mathbf{p}_1$ gives an extra contribution

$$\frac{e_1}{6m_1^2} \mathbf{x}_1 \times (\mathbf{s}_1 \times \mathbf{r}_{12}) \frac{\alpha_s}{r_{12}^3} \quad (8)$$

to $\boldsymbol{\mu}_1$. There are also possible orbital contributions to the moments because the Hamiltonian mixes states with nonzero orbital angular momenta with the ground state. These have the standard form to the accuracy we need.

The baryon moments are now given by expressions of the form

$$\mu = \sum_{i=1}^3 \mu_i \langle \sigma_{i,z} \rangle (1 + \delta_i) + \sum_{i=1}^3 \mu_i \langle L_{i,z} \rangle, \quad (9)$$

where we have quantized along $\mathbf{B}$, taken along the z axis. The expectation values are to be calculated in the baryon ground states. The correction terms δ_i from the new operators are given for the $L = 0$ baryons other than the Λ by

$$\begin{aligned} \delta_i &= \frac{3\epsilon_0 + \epsilon_1}{2m_1} + \frac{e_3}{e_1} \frac{\epsilon_2}{m_3} - \frac{\Delta_0 + \Delta_1}{2m_1}, \quad i = 1, 2, \\ \delta_3 &= \frac{\epsilon_2}{m_3} + \frac{2e_1}{e_3} \frac{\epsilon_1}{m_1} - \frac{\Delta_2}{m_3}, \end{aligned} \quad (10)$$

where the ϵ 's and Δ 's are ground state radial matrix elements,

$$\begin{aligned} \epsilon_0 &= \left\langle \frac{2}{3} \frac{\alpha_s}{6r_{12}} \right\rangle, & \epsilon_1 &= \left\langle \frac{2}{3} \frac{\alpha_s \mathbf{r}_{23} \cdot \mathbf{z}_2}{3r_{23}^3} \right\rangle, & \epsilon_2 &= \left\langle \frac{2}{3} \frac{\alpha_s \mathbf{r}_{31} \cdot \mathbf{z}_3}{3r_{31}^3} \right\rangle \\ \Delta_0 &= \left\langle \frac{\sigma r_{12}}{12} \right\rangle, & \Delta_1 &= \left\langle \frac{\sigma \mathbf{r}_{23} \cdot \mathbf{z}_2}{6r_{23}} \right\rangle, & \Delta_2 &= \left\langle \frac{\sigma \mathbf{r}_{31} \cdot \mathbf{z}_3}{6r_{31}} \right\rangle. \end{aligned} \quad (11)$$

The identical quarks in these baryons are labelled 1 and 2, the unlike quark, 3. In writing these results, we have made the approximation $r_1 + r_2 + r_3 \approx \frac{1}{2}(r_{12} + r_{23} + r_{31})$, known to be reasonably accurate for the ground state baryons [2], and used the corresponding Thomas spin interaction. The result for the Λ is similar.

2.3 Tests of the model

Rough estimates of the matrix elements above suggest that $\epsilon_i/m_l \approx 0.05$ and that $\Delta_i/m_l \approx 0.3$ for the light-quark masses used in Refs. 2 and 3. Binding effects are therefore potentially large, and are different for different quarks and baryons. To obtain a quantitative test of these effects and reliable estimates of the orbital contributions to the moments, expected to be small, we have performed a detailed analysis of the baryon wavefunctions using the Hamiltonian given in [1]. We use Jacobi-type internal coordinates $\boldsymbol{\rho}, \boldsymbol{\lambda}$, with $\boldsymbol{\rho} = \mathbf{x}_1 - \mathbf{x}_2$, and $\boldsymbol{\lambda} = \mathbf{R}_{12} - \mathbf{x}_3$, where $\mathbf{R}_{ij}$ is the coordinate of the center of mass of quarks i and j . The most general $j = \frac{1}{2}^+$ baryon wave function for quarks 1 and 2 identical and orbital angular momenta $L_\rho, L_\lambda, L \leq 2$ has the form

$$\begin{aligned} \psi_{\frac{1}{2},m} = & [a_0 \psi_a \mathbf{1} + i b_0 \psi_b (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \cdot \boldsymbol{\rho} \times \boldsymbol{\lambda} + c_0 \psi_c t_{12}(\boldsymbol{\rho}) \\ & + d_0 \psi_d t_{12}(\boldsymbol{\lambda}) + e_0 \psi_d (\boldsymbol{\sigma}_1 - \boldsymbol{\sigma}_2) \cdot \boldsymbol{\sigma}_3 \boldsymbol{\rho} \cdot \boldsymbol{\lambda}] \chi_{\frac{1}{2},m}^{S_{12}=1}. \end{aligned} \quad (12)$$

Here t_{12} is the usual tensor operator

$$t_{12}(\mathbf{x}) = 3\boldsymbol{\sigma}_1 \cdot \mathbf{x} \boldsymbol{\sigma}_2 \cdot \mathbf{x} - \boldsymbol{\sigma}_1 \cdot \boldsymbol{\sigma}_2 \mathbf{x}^2, \quad (13)$$

and $\chi_{\frac{1}{2},m}^{S_{12}=1}$ is the standard three-particle spinor for $S_{12} = 1$, $j = \frac{1}{2}$, $j_3 = m$. The scalar functions $\psi_i = \psi_i(\rho^2, \lambda^2)$ are normalized together with the accompanying spin operators. The constant coefficients $a_0, \dots, e_0$, normalized to unity, give the fractions of the various component states in ψ . With this form of the wave function, we can use trace methods for the spins to calculate such quantities as $\langle H \rangle$ and $\langle \boldsymbol{\mu} \rangle$.

We have made variational calculations of the energies and approximate wave functions of the ground state baryons and their first excited states using the Hamiltonian in [1]. The information on the excited states allows us to calculate the coefficients $b_0, \dots, e_0$ perturbatively. While the Hamiltonian does mix orbitally excited states into the $L_\rho = L_\lambda = L = 0$ quark-model ground state, the coefficients are very small, ranging from essentially zero to about 0.02 depending on the baryon. Because these coefficients only appear quadratically in the baryon moments, the orbital contributions to the moments are completely negligible.

The radial matrix elements ϵ_i and Δ_i defined in Eq. (11) are significant, with the ϵ 's ranging from 10 to 20 MeV, and the Δ 's from 40 to 70 MeV. A new fit to the moment data using the expressions in Eqs. (9) and (10), with the quark

masses allowed to vary, gives a small improvement in the fitted moments, with the root-mean-square deviation from the measured moments decreasing from 0.14 nm for the quark model to 0.10 nm. A secondary effect of the corrections is to change the fitted quark masses or moments significantly, with the effective quark masses decreasing, or the moments increasing.

We have concluded from this exercise first, that the finer details of baryon structure are not yet described correctly in the present QCD-based model, and second, that the baryon magnetic moments provide a very sensitive check on the theory.

3 Possible improvements in the theory

We begin the discussion of possible improvements in the theory of the moments by recalling the approximations used in the construction given by Brambilla *et al.* [1] and in our derivation of the moment operators:

- (i) the $1/m_q$ expansion;
- (ii) the minimal surface approximation;
- (iii) forward propagation of the quarks in time;
- (iv) and the quenched approximation.

Of these approximations, only (iii) and (iv) are likely to affect the moments significantly.

The $1/m_q$ expansion is to be interpreted as an expansion in constituent quark masses. It can be resummed in the kinetic terms in the Hamiltonian as in Eq. (4), and leads elsewhere to the appearance of effective inverse masses $1/m_i$ which need not be the same as the kinematic masses, but represent averages of quantities such as $1/E_i = 1/\sqrt{p_i^2 + m_i^2}$. Since there is no new spin dependence involved, and we have treated the quark masses as free parameters in fitting the moments, relativistic corrections of this sort are unlikely to change our results.

The average over the color gauge field A_g in the minimal surface approximation neglects fluctuations, and minimizes the surface energy of the world sheet to obtain the approximate Hamiltonian. Since the color fields do not carry charge, they do not contribute internal currents in the baryon, and an improved treatment of the averaging would presumably not change the moment operator directly. While it could change the functional form of the Hamiltonian somewhat, the quantitative success of the model in describing baryon spectra suggests that the changes would not be large, and their effect on the moments through the ϵ and Δ parameters would be minimal.

In contrast, the remaining two approximations affect the moment operator directly. Internal quark loops can contribute circulating currents in the baryon. These are omitted in the quenched approximation. In addition, the forward

propagation of the quarks in time inherent in the use of the Foldy-Wouthuysen approximation eliminates quark pair effects connected with the valence lines in Fig. 1. We believe that this general quenched picture underlies the difficulties with the model, and that it will be necessary to include pair effects to obtain a precise dynamical description of the moments.

One effect of internal quark loops can be seen in Fig. 2. In this figure, we suppose that a quark loop is embedded in the minimal world sheet of the baryon. The effect is to give a meson state and a new baryon in a world-sheet picture of meson emission and absorption. An external magnetic field interacting with the




Figure 2: Quark loops embedded in the world sheet give meson-baryon states.

system will see a meson current as well as the baryon current, and the moment will be modified. Meson currents were invoked in the past in attempts to explain the *full* anomalous moments of the nucleons. Here we are only concerned with presumably small corrections to the quark-model moments.

One approach to the calculation of meson loop effects is through chiral perturbation theory. This has a long history, and has been studied recently in the context of baryon moments by a number of authors [4, 5, 6]. The relevant diagrams for the interaction of the baryons with the pseudo Goldstone bosons of the chiral theory are shown in Fig. 3. The work of Jenkins *et al.* [4] uses




Figure 3: Goldstone-boson diagrams which contribution to the magnetic moments.

heavy baryon chiral perturbation theory [7], and studies the effect on the moments of the “nonanalytic terms” in the symmetry breaking parameter m_s . The




Figure 4: (a) Appearance of Bethe-Salpeter wave functions in the diagrams. (b) Schematic reminder of the distributed nature of the meson-baryon vertex.

analytic terms in m_s are ambiguous because the inevitable appearance of new couplings at each order in the chiral expansion, and are ignored. The results of Jenkins *et al.* are not encouraging as they stand, but we have found errors in some of the coupling factors given in the published paper. Luty *et al.* [5] concentrate on a simultaneous expansion in $1/N_c$ and m_s , and obtain interesting sum rules for the moments but little dynamical information. Finally, Bos *et al.* [6] consider the moments from the point of view of flavor $SU(3)$ breaking in the baryon octet using a different chiral counting than that usually used, and obtain a very successful parametrization for the moments. However, this model is again nondynamical, and has seven parameters to describe eight measured moments. In fact, a fundamental problem with the chiral expansion from our perspective is that it simply parametrizes the moments with an expansion consistent with QCD, but does not control the higher order terms in what appears to be a slowly convergent series.

We are presently investigating meson loop effects using a somewhat different approach suggested by the world-sheet picture. The baryon appears in this picture as an extended object which must absorb the recoil momentum in the emission of a meson. We would therefore expect wave function effects (form factors) to be important for high meson momenta, and to supply a natural cutoff for loop graphs. The wave function appears naturally when the process is viewed using “old fashioned” instead of Feynman perturbation theory, as indicated in Fig. 4: energy denominators and vertex functions combine to give the $B'M$ component of the wave function for a baryon B , or, with different time ordering, the BM component of the wave function of B' . Since the wave functions are expected to be fairly soft, with characteristic momenta below the chiral cutoff of ~ 1 GeV, they can be expected to combine a number of terms in the chiral expansion in a way that is dynamically accessible through approximate models derived from QCD such as that of Brambilla *et al.* [1]. The non-point character of the meson-baryon vertex in spacetime indicated in Fig. 4 contrasts sharply with the point vertex used in chiral perturbation theory, Fig. 3, and is

also likely to play a role. We have reached the same conclusions by studying the exact sideways dispersion relations for the moments given by Bincer [8]. The problem there is in extracting the quark-model moments.

We note finally that a different aspect of symmetry breaking, the suppression of strange-quark loops through mass effects, appears as a natural possibility in a world-sheet picture. Note that this is *not* the same as the suppression of kaon loops considered by other authors; see, e.g., [4] and the references given there.

4 Conclusions

On the basis of the work sketched above, we have concluded that the magnetic moments of the baryons give a sensitive test of baryon structure and of approximations in QCD. In particular, the quenched approximation, while reasonably successful when used in the calculation of baryon and meson spectra, appears to fail for the moments. The accurate calculation of the moments by lattice methods will presumably involve going beyond that approximation, and will provide a precision test of the methods used.

We find also that the world-sheet picture of baryon structure gives useful insights into the moments problem, and provides a new point of view which could be developed further. Problems which need further study within this approach to QCD include the following:

- (i) establishing the connection to, and the relevance of, the chiral limit;
- (ii) the development of methods to incorporate loop effects which build in the extended structure of the baryons;
- (iii) and the possible usefulness of string theory methods in the calculation of loop effects.

Acknowledgments

The authors have benefited from conversations with Dr. Nora Brambilla, and appreciate her interest in this work, and her organization of the Como Conference. This research was supported in part by the U.S. Department of Energy under Grant No. DE-FG02-95ER40896, and in part by the University of Wisconsin Graduate School with funds from the Wisconsin Alumni Research Foundation.

References

References

- [1] N. Brambilla, P. Consoli, and G.M. Prosperi, Phys. Rev. D **50**, 5878 (1994).

- [2] J. Carlson, J. Kogut, and V.R. Pandharipande, Phys. Rev. D **27**, 233 (1983); **28**, 2807 (1983).
- [3] S. Capstick and N. Isgur, Phys. Rev. D **34**, 2809 (1986).
- [4] E. Jenkins, M. Luke, A.V. Manohar, and M.J. Savage, Phys. Lett. B **302**, 482 (1993).
- [5] M.A. Luty, J. March-Russell, and M. White, Phys. Rev. D **51**, 2332 (1995).
- [6] J.W. Bos, D. Chang, S.C. Lee, Y.C. Lin, and H.H. Shih, hep-ph/9602251.
- [7] E. Jenkins and A.V. Manohar, Phys. Lett. B **255**, 558 (1991).
- [8] A. Bincer, Phys. Rev. **118**, 855 (1960).